arXiv:0709.3196v1 [quant-ph] 20 Sep 2007

Quantum nonlocality without entanglement in a pair of qubits

Masato Koashi, Fumitaka Takenaga, Takashi Yamamoto, and Nobuyuki Imoto
Division of Materials Physics, Department of Materials Engineering Science,
Graduate School of Engineering Science, Osaka University,
1-3 Machikaneyama, Toyonaka, Osaka 560-8531, Japan and
CREST Photonic Quantum Information Project,
4-1-8 Honmachi, Kawaguchi, Saitama 331-0012, Japan

We consider unambiguous discrimination of two separable bipartite states, one being pure and the other being a rank-2 mixed state. There is a gap between the optimal success probability under global measurements and the one achieved by generalized measurements with separable measurement operators. We show that even the latter success probability cannot be achieved via local operations and classical communication, leaving a nonzero gap in between.

In the recent progress on understanding of quantum information, distinction between classical interaction and quantum interaction has played a key role. For a bipartite system AB, a restricted class of operations called LOCC (local operations and classical communication) is considered, and any state that can be prepared from scratch under LOCC is called separable. The rest of the states are termed entangled states, which are prepared only through quantum interaction between A and B. One might expect that the classical/quantum boundary would be clear-cut in this paradigm, and separable states would have nothing to do with the quantum interaction. But that is not the case. Peres and Wootters suggested [1] that a set of bipartite separable states can be distinguished better by a global measurement, which can be realized only by quantum interaction between the two systems. Later, using different sets of states and different measures of distinguishability, a stronger evidence was shown [2], a nonzero gap between the optimal global distinguishability $P_{\rm opt}^{\rm (glo)}$ and the LOCC counterpart $P_{\rm opt}^{\rm (LOCC)}$ was found [3], and more recently, global measurements surpassing theoretically derived $P_{\rm opt}^{\rm (LOCC)}$ were experimentally demonstrated [4, 5].

If a measurement is implementable under LOCC, it must be a *separable measurement*, for which every POVM (positive-operator-valued measure) element is separable (written as a sum of positive product operators like $\sum_j \hat{A}_j \otimes \hat{B}_j$). In most of the above examples, the measurement achieving $P_{\rm opt}^{\rm (glo)}$ is inseparable, and hence one can use it to create entanglement from a separable state [6]. The reason why the measurement cannot be carried out under LOCC is obvious in this respect. An exception is the example given in [3], composed of a cleverly chosen set of nine (or eight) pure orthogonal states of a pair of qutrits (three level systems). In this case, the best result is achieved by a separable measurement. Nonetheless, the LOCC measurements are proved to be less efficient and there is a nonzero gap between $P_{\rm opt}^{\rm (sep)}$ and $P_{\rm opt}^{\rm (LOCC)}$, where $P_{\rm opt}^{\rm (sep)}$ is the optimal distinguishability among separable measurements. This implies that the optimal separable measurement has a striking irreversibility under LOCC: we must invest entanglement to implement the measurement, whereas we can never use it to create entanglement from scratch. We see interesting parallels between this and a bound entangled state [7]. For the latter, we must invest entanglement to create the state, whereas we can never extract from it a standard form of entanglement like a singlet state of two spin-1/2 particles. In comparison to bound entanglement, our understanding about such bizarre measurements or processes are still rudimentary, including the very reason why the LOCC cannot implement those. For example, we know that the bound entanglement is unique to higher-level systems and does not appear in a qubit-pair system [8], but we know little about the same question for $P_{\rm opt}^{\rm (sep)} > P_{\rm opt}^{\rm (LOCC)}$.

The aim of this paper is to provide a discrimination problem exhibiting such a phenomenon, which is much simpler in many ways: the system is a qubit pair, and the number of separable states to distinguish is just two. The description of the two states involve only three pure product states. The two states may look rather mundane and with little contraption, but we can still prove $P_{\rm opt}^{\rm (glo)} > P_{\rm opt}^{\rm (sep)} > P_{\rm opt}^{\rm (LOCC)}$. The key element in the proof is the observation that refinement process of any LOCC measurement must proceed alternately between the two systems. This should be contrasted to the property used in the proof of [3], that the refinement process can always be regarded as a continuous one.

The problem we consider here is the unambiguous discrimination [9] between the two separable states of a two-qubit system defined by

$$\hat{\rho}_0 = |00\rangle\langle 00|, \tag{1}$$

$$\hat{\rho}_1 = (1/2)(|++\rangle\langle ++| + |--\rangle\langle --|), \tag{2}$$

where $|\pm\rangle \equiv 2^{-1/2}(|0\rangle \pm |1\rangle)$. The two qubits are secretly prepared in one of the two states before one qubit is sent to Alice and the other one to Bob. They should determine the identity of the prepared state by a quantum measurement. We allow them to declare that the measurement has been a failure, but otherwise they must identify the state with no errors. Such a measurement is generally described by a POVM $\{\hat{F}_m\}_{m=0,1,2}$ satisfying

$$\mathrm{Tr}(\hat{F}_0\hat{\rho}_1) = \mathrm{Tr}(\hat{F}_1\hat{\rho}_0) = 0, \tag{3}$$

where the outcome $m = 2$ corresponds to the failure and $m = 0, 1$ to the identification of state $\hat{\rho}_m$. For $j = 0, 1$, let $\gamma_j \equiv \mathrm{Tr}(\hat{F}_j\hat{\rho}_j)$ be the probability of success when the prepared state was $\hat{\rho}_j$. What we seek is the supremum of γ_1 over all the allowed measurements with a fixed value of γ_0, which we will denote by $P_{\rm opt}(\gamma_0)$. A related problem is the optimization of the averaged success rate $\gamma_{\rm ave} \equiv \eta_0\gamma_0 + \eta_1\gamma_1$ when ρ_j is prepared with probability η_j. The optimized value $Q_{\rm opt}$ can be straightforwardly calculated once we learn the function $P_{\rm opt}(\gamma_0)$, namely, $Q_{\rm opt} = \max_\gamma[\eta_0\gamma + \eta_1 P_{\rm opt}(\gamma)]$.

If we allow Alice and Bob arbitrary global quantum operations, the optimal probability $\gamma_1 = P^{(\rm glo)}_{\rm opt}(\gamma_0)$ is known [10] to be achieved by the following protocol. First, they globally measure whether their bit values coincide, namely, project the state to the subspace spanned by $\{|00\rangle, |11\rangle\}$ and to the one by $\{|01\rangle, |10\rangle\}$. If the prepared state was $\hat{\rho}_1$, the latter case occurs with probability 1/2 and then the measurement successfully identifies it. Otherwise, they are left with the discrimination between $|\psi_0\rangle \equiv |00\rangle$ and $|\psi_1\rangle \equiv 2^{-1/2}(|00\rangle + |11\rangle)$ with $s \equiv |\langle\psi_0|\psi_1\rangle|^2 = 1/2$. This is a well-known problem and $|\psi_0\rangle$ can be identified with probability $\gamma_0(\leq 1/2)$ while $|\psi_1\rangle$ can be identified with $1 - s/(1-\gamma_0)$ [11]. The whole protocol achieves γ_1 equal to

$$P^{(\rm glo)}_{\rm opt}(\gamma_0) = 1 - [4(1-\gamma_0)]^{-1} \tag{4}$$

for $0 \leq \gamma_0 \leq 1/2$, which is plotted in Fig. 1 (a).



FIG. 1: (a) Bounds on success probabilities (γ_0, γ_1). (b) Expanded after taking the dotted straight line in (a) as the base line. $P^{(\rm LOCC)}_{\rm opt}$ should lie in the shaded region.

Next, we determine the optimal probability $\gamma_1 = P^{(\rm sep)}_{\rm opt}(\gamma_0)$ among separable measurements. It is convenient to use the high symmetry of states $\hat{\rho}_0$ and $\hat{\rho}_1$. These states are invariant under any of the following maps — Exchange of the two qubits, given by the map $S^{(1)} : |ij\rangle\langle kl| \mapsto |ji\rangle\langle lk|$; simultaneous phase flip, given by $S^{(2)} : |ij\rangle\langle kl| \mapsto (-1)^{i+j+k+l}|ij\rangle\langle kl|$; (global) transpose, given by $S^{(3)} : |ij\rangle\langle kl| \mapsto |kl\rangle\langle ij|$; and partial transpose, given by $S^{(4)} : |ij\rangle\langle kl| \mapsto |il\rangle\langle kj|$. As a result, if a separable measurement $\{\hat{F}_m\}$ achieves success probabilities (γ_0, γ_1), all the POVMs generated by applying the above maps are physically valid (note the separability of $\hat{F}_m$ for $S^{(4)}$ [12]) and give the same probabilities (γ_0, γ_1). Then, the symmetrized measurement, which executes one of those 2^4 measurements randomly, also achieves the same (γ_0, γ_1). Let us define the symmetrizing map via the composition $\mathcal{S} \equiv \prod_{n=1}^4 2^{-4}(I + S^{(n)})$, where I is the identity map. The POVM of the symmetrized measurement is given by $\{\mathcal{S}(\hat{F}_m)\}$. Any symmetrized self-adjoint operator has a simple form with four real parameters in the matrix representation on the basis $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$ as

$$[a, b, c; \mu] \equiv \begin{pmatrix} a & 0 & 0 & \mu \\ 0 & c & \mu & 0 \\ 0 & \mu & c & 0 \\ \mu & 0 & 0 & b \end{pmatrix}. \tag{5}$$

It is positive (semidefinite) if and only if $a \geq 0$, $ab \geq \mu^2$, and $c \geq |\mu|$. With this notation, the two initial states are written by $\hat{\rho}_0 = [1, 0, 0; 0]$ and $\hat{\rho}_1 = (1/4)[1, 1, 1; 1]$.

The conditions $\mathrm{Tr}[\hat{\rho}_0\mathcal{S}(\hat{F}_0)] = \gamma_0$ and $\mathrm{Tr}[\hat{\rho}_1\mathcal{S}(\hat{F}_0)] = 0$ require that $\mathcal{S}(\hat{F}_0) = \gamma_0[1, 1, 1; -1]$. To satisfy $\mathrm{Tr}[\hat{\rho}_0\mathcal{S}(\hat{F}_1)] = 0$ and to be positive, $\mathcal{S}(\hat{F}_1)$ must be written as $\mathcal{S}(\hat{F}_1) = [0, b, c; 0]$. The positivity of $\mathcal{S}(\hat{F}_2) = [1-\gamma_0, 1-\gamma_0-b, 1-\gamma_0-c; \gamma_0]$ requires $c \leq 1 - 2\gamma_0$ and $b \leq (1-2\gamma_0)/(1-\gamma_0)$. The optimal value of $\gamma_1 = \mathrm{Tr}[\hat{\rho}_1\mathcal{S}(\hat{F}_1)] = (b + 2c)/4$ is thus given by

$$P^{(\rm sep)}_{\rm opt}(\gamma_0) = 1 - \gamma_0 - [4(1-\gamma_0)]^{-1} \tag{6}$$

for $0 \leq \gamma_0 \leq 1/2$, which is shown in Fig. 1. This is achieved if and only if $\mathcal{S}(\hat{F}_m) = \hat{F}^{(\rm sep)}_m$ with

$$\begin{aligned} \hat{F}^{(\rm sep)}_0 &\equiv 2\gamma_0(\hat{P}_+ \otimes \hat{P}_- + \hat{P}_- \otimes \hat{P}_+) & (7)\\ \hat{F}^{(\rm sep)}_1 &\equiv (1-2\gamma_0)(\hat{P}_0 \otimes \hat{P}_1 + \hat{P}_0 \otimes \hat{P}_1) & \\ &\quad +(1-\xi_0)\hat{P}_1 \otimes \hat{P}_1 & (8)\\ \hat{F}^{(\rm sep)}_2 &\equiv [(1+\xi_0)/2](\hat{P}_{\gamma+} \otimes \hat{P}_{\gamma+} + \hat{P}_{\gamma-} \otimes \hat{P}_{\gamma-}) & (9) \end{aligned}$$

where $\xi_0 \equiv \gamma_0/(1-\gamma_0)$, $|\gamma\pm\rangle \equiv \sqrt{1-\gamma_0}|0\rangle \pm \sqrt{\gamma_0}|1\rangle$, and we denoted $\hat{P}_X \equiv |X\rangle\langle X|$.

Next, for $0 < \gamma_0 < 1/2$, we will show that the optimal separable measurement $\{\hat{F}^{(\rm sep)}_m\}$ cannot be implemented under LOCC, and there is a nonzero gap between $P^{(\rm sep)}_{\rm opt}$ and $P^{(\rm LOCC)}_{\rm opt}$. In a general LOCC measurement protocol, Alice first obtains an outcome i_1 and tells Bob. If they stopped the protocol here, it would be regarded as a measurement with POVM $\{\hat{G}_{i_1}\}_{i_1}$ with $\hat{G}_{i_1} = \hat{A}_{i_1} \otimes \hat{1}$. Next, Bob obtains an outcome i_2 and tells Alice. The POVM is refined to $\{\hat{G}_{i_1 i_2}\}_{i_1,i_2}$ with $\hat{G}_{i_1 i_2} = \hat{A}_{i_1} \otimes \hat{B}_{i_1 i_2}$ at this



FIG. 2: Refining process of an LOCC measurement.

point, where $\sum_{i_2} \hat{G}_{i_1 i_2} = \hat{G}_{i_1}$. Repeating such a process n rounds, they carry out a measurement with POVM $\{\hat{G}_{i_1 \ldots i_n}\}_{i_1,\ldots,i_n}$, with $\hat{G}_{i_1 \ldots i_n} = \hat{A}_{i_1 i_2 \cdots} \otimes \hat{B}_{i_1 i_2 \cdots}$. Finally, they determine the final outcome $m = 0, 1, 2$ according to a rule $m = \Omega(i_1, \ldots, i_n)$.

Let us introduce a "weight" of a POVM element $\hat{G}$ by $w(\hat{G}) \equiv \langle 00|\hat{G}|00\rangle = \mathrm{Tr}(\hat{\rho}_0 \hat{G})$. As long as $w > 0$, we can keep track of the progress $\hat{1} \otimes \hat{1} \to \hat{G}_{i_1} \to \hat{G}_{i_1 i_2} \to \cdots \to \hat{G}_{i_1 \ldots i_n}$ using the two real parameters defined by

$$x(\hat{A}) \equiv \mathrm{Re}\, A_{01}/A_{00},\ y(\hat{B}) \equiv \mathrm{Re}\, B_{01}/B_{00}, \tag{10}$$

where $A_{jk} \equiv \langle j|\hat{A}|k\rangle$ and $B_{jk} \equiv \langle j|\hat{B}|k\rangle$. The starting point for $\hat{1} \otimes \hat{1}$ is $(x(\hat{1}), y(\hat{1})) = (0, 0)$. This is followed by the point $(x(\hat{A}_{i_1}), 0)$, representing $\hat{G}_{i_1}$, and then the point $(x(\hat{A}_{i_1}), y(\hat{B}_{i_1 i_2}))$ for $\hat{G}_{i_1 i_2}$, and so on. We notice that either x or y, and not both, changes in a single step. Hence the trajectory on the xy plane is a zigzag line as in Fig. 2. This property, reflecting the fact that Alice and Bob can refine the measurement operator *only alternately* to arrive at the final outcome, is crucial in showing that the optimal separable measurement is not feasible.

The parameters w, x, y take distinct values for $m = 0, 1$. If $m = \Omega(i_1, \ldots, i_n) = 0$ and hence $\mathrm{Tr}(\hat{G}_{i_i \ldots i_n} \hat{\rho}_1) = 0$, then $\hat{G}_{i_1 \ldots i_n}$ should be proportional to $\hat{P}_+ \otimes \hat{P}_-$ or to $\hat{P}_- \otimes \hat{P}_+$, namely,

$$(x, y) = (1, -1) \ \text{or} \ (-1, 1) \ \text{for} \ m = 0. \tag{11}$$

If $m = \Omega(i_1, \ldots, i_n) = 1$, $\mathrm{Tr}(\hat{G}_{i_i \ldots i_n} \hat{\rho}_0) = 0$ means that

$$w(\hat{G}_{i_i \ldots i_n}) = 0 \ \text{for} \ m = 1. \tag{12}$$

In order to achieve $\gamma_1 = P_{\mathrm{opt}}^{(\mathrm{sep})}(\gamma_0)$, the outcomes with $m = \Omega(i_1, \ldots, i_n) = 2$ must also have distinct values for (x, y), namely,

$$(x, y) = (\sqrt{\xi_0}, \sqrt{\xi_0}) \ \text{or} \ (-\sqrt{\xi_0}, -\sqrt{\xi_0}) \ \text{for} \ m = 2. \tag{13}$$

This is because we see from Eq. (9) that the range of $\mathcal{S}(\hat{F}_2) = \hat{F}_2^{(\mathrm{sep})}$ is spanned by $\{|\gamma+\rangle|\gamma+\rangle, |\gamma-\rangle|\gamma-\rangle\}$ and includes no other product states since $\gamma_0 > 0$. Hence $\hat{G}_{i_1 \ldots i_n}$ should be proportional to $\hat{P}_{\gamma\pm} \otimes \hat{P}_{\gamma\pm}$. This allows us to find a pair of linear functionals $f_\pm$ that are nonpositive for any final outcome $\hat{G}_{i_1 \ldots i_n}$. These are defined by

$$\begin{aligned} f_\pm(\hat{F}) \equiv\ & F_{0011} + F_{0110} + F_{1001} + F_{1100} - 4\xi_0 F_{0000} \\ & \pm (1 + \xi_0)(F_{0010} + F_{1000} - F_{0001} - F_{0100}), \end{aligned} \tag{14}$$

where $F_{ijkl} \equiv \langle ij|\hat{F}|kl\rangle$. For a product operator $\hat{G} = \hat{A} \otimes \hat{B}$ with $x(\hat{A}) = x$ and $y(\hat{B}) = y$, we have

$$\begin{aligned} f_\pm(\hat{G}) &= w(\hat{G})[4(xy - \xi_0) \pm 2(1 + \xi_0)(x - y)] \quad (15) \\ &= w(\hat{G})[\chi_\pm(x, y) + (1 - \xi_0)^2], \quad (16) \end{aligned}$$

where $\chi_\pm(x, y) \equiv [2x \mp (1 + \xi_0)][2y \pm (1 + \xi_0)]$. Comparing Eqs. (11)–(13) and (15), we conclude that $f_\pm(\hat{G}_{i_1 \ldots i_n})$ should be either 0 or $-8(1 + \xi_0)w$ in order to achieve $\gamma_1 = P_{\mathrm{opt}}^{(\mathrm{sep})}(\gamma_0)$. As we will see, this is impossible with LOCC.

Since $\gamma_0 > 0$, there is an outcome with $m = 0$ satisfying $w(\hat{G}_{i_1 \cdots i_n}) > 0$. From Eq. (11), we may assume that $(x, y) = (1, -1)$ for this outcome (since the case for $(-1, 1)$ is similar). Let us define regions $R_\pm \equiv \{(x, y)|\chi_\pm(x, y) \geq 0\}$, which are shown by shaded areas in Fig. 2. It is obvious that the zigzag path from the origin must land on R_+ at least once in order to reach $(1, -1)$. Let $\hat{G}_{i_1 \cdots i_m}$ be the point on R_+. From Eq. (16), we see $f_+(\hat{G}_{i_1 \cdots i_m}) \geq w(\hat{G}_{i_1 \cdots i_m})(1 - \xi_0)^2 \geq w(\hat{G}_{i_1 \cdots i_n})(1 - \xi_0)^2 > 0$ since $\gamma_0 < 1/2$. This implies that there exists at least one final outcome satisfying $f_+(\hat{G}_{i_1 \cdots i_m j_{m+1} \cdots j_n}) > 0$. Hence we conclude that no finite-round LOCC protocols achieve $\gamma_1 = P_{\mathrm{opt}}^{(\mathrm{sep})}(\gamma_0)$.

Since the above reasoning still allows for the existence of LOCC protocols achieving γ_1 arbitrarily close to $P_{\mathrm{opt}}^{(\mathrm{sep})}(\gamma_0)$, we need a more quantitative argument to show that there is a nonzero gap $P_{\mathrm{opt}}^{(\mathrm{sep})}(\gamma_0) - P_{\mathrm{opt}}^{(\mathrm{LOCC})}(\gamma_0) > 0$. First, we divide all the possible final outcomes $(i_1, \ldots, i_n)$ into three groups $\Gamma_0, \Gamma_+, \Gamma_-$ according to the trajectories $\hat{1} \otimes \hat{1} \to \hat{G}_{i_1} \to \hat{G}_{i_1 i_2} \to \cdots \to \hat{G}_{i_1 \ldots i_n}$ by the following rules. (i) $(i_1, \ldots, i_n) \in \Gamma_0$ if the trajectory never goes into the region $R_+ \cup R_-$ as long as the weight w is positive. (ii) $(i_1, \ldots, i_n) \in \Gamma_+$ if the first point of trajectory in the region $R_+ \cup R_-$ is in the region R_+. (iii) $(i_1, \ldots, i_n) \in \Gamma_-$ otherwise. It is obvious that in the last case, the first point of trajectory in the region $R_+ \cup R_-$ is in R_-.

Let $\hat{K}_m^{(\pm)}$ be the sum of the final POVM elements $\hat{G}_{i_1 \ldots i_n}$ satisfying $(i_1, \ldots, i_n) \in \Gamma_\pm$ and $\Omega(i_1, \ldots, i_n) = m$. Define $\hat{K}^{(\pm)} = \hat{K}_0^{(\pm)} + \hat{K}_1^{(\pm)} + \hat{K}_2^{(\pm)}$. Since $\hat{K}^{(+)}$ is also written as the sum of operators $\hat{G}_{i_1 \ldots i_m}$ for the first points entering R_+, we have $f_+(\hat{K}^{(+)}) \geq w(\hat{K}^{(+)})(1 - \xi_0)^2$ from Eq. (16). On the other hand, we see $f_+(\hat{K}_0^{(+)}) \leq 0$ and $w(\hat{K}_1^{(+)}) = f_+(\hat{K}_1^{(+)}) = 0$ from Eqs. (11), (12), and (15). Then we have

$$f_\pm(\hat{K}_2^{(\pm)}) \geq [w(\hat{K}_0^{(\pm)}) + w(\hat{K}_2^{(\pm)})](1 - \xi_0)^2, \tag{17}$$

where a similar result for Γ_- is included. As we have seen before, no outcome for $m = 0$ belongs to Γ_0. Hence $\hat{K}_0^{(+)} + \hat{K}_0^{(-)} = \hat{F}_0$ and

$$w(\hat{K}_0^{(+)}) + w(\hat{K}_0^{(-)}) = \gamma_0. \tag{18}$$

The next step is to quantify how much penalty is imposed on γ_1 when Eq. (13) is not satisfied. Let $\hat{N}$ be a positive separable operator. We will prove that if $\hat{F}_2 - \hat{N}$ is positive, $\gamma_1 \leq P_{\rm opt}^{(\rm sep)}(\gamma_0) - g(\hat{N})$ where $g(\hat{N}) \equiv (1/4)\mathrm{Tr}\{\hat{N}[\xi_0^2, 1, 1; -(1+\xi_0)/2]\}$.

Let $\mathcal{S}(\hat{N}) = [a', b', c'; \mu']$ and $\mathcal{S}(\hat{F}_1) = \hat{F}_1^{(\rm sep)} - [0, \Delta b, \Delta c; 0]$. $\mathcal{S}(\hat{F}_2 - \hat{N}) = \hat{F}_2^{(\rm sep)} - \mathcal{S}(\hat{N}) + [0, \Delta b, \Delta c; 0]$ $= [1 - \gamma_0 - a', \gamma_0\xi_0 - b' + \Delta b, \gamma_0 - c' + \Delta c; \gamma_0 - \mu']$ must be positive. This leads to (i) $\Delta c \geq c' - \mu'$, which is obvious, and (ii) $\Delta b \geq \xi_0^2 a' + b' - 2\xi_0\mu'$, which is proved as follows. When $a' = 1 - \gamma_0$, $\mu' = \gamma_0$ must hold and then (ii) is true since $\gamma_0\xi_0 - b' + \Delta b \geq 0$. When $a' < 1 - \gamma_0$, assuming that (ii) is false would lead to

$$\begin{aligned}(1 - \gamma_0 - a')(\gamma_0\xi_0 - b' + \Delta b) - (\gamma_0 - \mu')^2 \\ < -(\xi_0 a' - \mu')^2 \leq 0,\end{aligned} \tag{19}$$

and hence (ii) must be true. Now from (i) and (ii), we have $P_{\rm opt}^{(\rm sep)}(\gamma_0) - \gamma_1 = (\Delta b + 2\Delta c)/4 \geq g(\hat{N})$.

For a positive product operator $\hat{G} = \hat{A} \otimes \hat{B}$ with $x(\hat{A}) = x$ and $y(\hat{B}) = y$, we can bound g by using the requirement for the positivity, $A_{11} \geq x^2 A_{00}$ and $B_{11} \geq y^2 B_{00}$, as

$$g(\hat{G}) \geq w(\hat{G})[(xy - \xi_0)^2 + (x - y)^2]/4. \tag{20}$$

Comparing it with Eq. (15), we have $\eta_0 w(\hat{G}) g(\hat{G}) \geq f_\pm(\hat{G})^2$ with $\eta_0 \equiv 16[4 + (1 + \xi_0)^2]$. This relation can be extended to any separable operator $\hat{K} = \sum_k \hat{G}_k$. In fact, with $p_k \equiv w(\hat{G}_k)/w(\hat{K})$, we have $\eta_0 w(\hat{K}) g(\hat{K}) = \sum_k \eta_0 w(\hat{G}_k) g(\hat{G}_k)/p_k \geq \sum_k p_k [f_\pm(\hat{G}_k)/p_k]^2 \geq [\sum_k f_\pm(\hat{G}_k)]^2 = f_\pm(\hat{K})^2$. Combining this with Eq. (17), we have

$$\begin{aligned} g(\hat{K}_2^{(\pm)}) &\geq \frac{[w(\hat{K}_0^{(\pm)}) + w(\hat{K}_2^{(\pm)})]^2(1 - \xi_0)^4}{\eta_0 w(\hat{K}_2^{(\pm)})} \\ &\geq 4\eta_0^{-1}(1 - \xi_0)^4 w(\hat{K}_0^{(\pm)}) \end{aligned} \tag{21}$$

Since $\hat{F}_2 - (\hat{K}_2^{(+)} + \hat{K}_2^{(-)})$ is positive, $\gamma_1 \leq P_{\rm opt}^{(\rm sep)}(\gamma_0) - g(\hat{K}_2^{(+)} + \hat{K}_2^{(-)})$. Hence, using Eq. (18), we have a bound $\gamma_1 \leq P_{\rm opt}^{(\rm sep)}(\gamma_0) - 4\eta_0^{-1}(1 - \xi_0)^4\gamma_0$ under LOCC, namely,

$$\begin{aligned} &P_{\rm opt}^{(\rm LOCC)}(\gamma_0) \leq u(\gamma_0) \\ &\equiv P_{\rm opt}^{(\rm sep)}(\gamma_0) - \frac{(1 - 2\gamma_0)^4\gamma_0}{4(1 - \gamma_0)^2[1 + 4(1 - \gamma_0)^2]}, \end{aligned} \tag{22}$$

which proves that there is a nonzero gap except $\gamma_0 = 0, 1/2$.

We also give lower bounds on $P_{\rm opt}^{(\rm LOCC)}(\gamma_0)$ by finding specific LOCC protocols. If Alice first measures on $\{|+\rangle, |-\rangle\}$, Bob's task is either the discrimination of $\{|0\rangle, |+\rangle\}$ or that of $\{|0\rangle, |-\rangle\}$, and hence they can achieve $\gamma_1 = l_1(\gamma_0) \equiv 1 - [2(1 - \gamma_0)]^{-1}$. If Alice and Bob both measure on $\{|0\rangle, |1\rangle\}$, they achieve $(\gamma_0, \gamma_1) = (0, 3/4)$. Mixture of the $(\gamma_0, \gamma_1) = (0, 3/4)$ protocol and the $(\gamma_0, \gamma_1) = (\sqrt{2} - 1, l_1(\sqrt{2} - 1))$ protocol gives another linear lower bound $l_2(\gamma_0)$. These are shown in Fig. 1 (b).

To summarize, we have found a simple problem of state discrimination with $P_{\rm opt}^{(\rm glo)} > P_{\rm opt}^{(\rm sep)} > P_{\rm opt}^{(\rm LOCC)}$. This means that, even for a qubit pair, there exists a measurement that requires entanglement to execute, but never produces entanglement from scratch. The measurement is still useful, having a definite advantage over any LOCC measurement. The pair of states in our example are very simple — in fact, they are one of the simplest in a sense, considering that $P_{\rm opt}^{(\rm glo)} = P_{\rm opt}^{(\rm LOCC)}$ has already been proved for discrimination of any two pure product states [13]. This suggests that the existence of the gap may not be rare but could show up in many other problems, all the more reason to investigate this striking irreversibility in more detail. The qubit pair is a convenient playground for such investigation, because operations are parametrized by fewer parameters and entanglement is understood more quantitatively. We hope that the present example would serve as a driving force toward a better understanding about the subtle boundary between quantum and classical interactions.

The authors thank K. Azuma for helpful discussions. This work was supported by a MEXT Grant-in-Aid for Young Scientists (B) 17740265.

[1] A. Peres and W. K. Wootters, Phys. Rev. Lett. **66**, 1119 (1991).

[2] S. Massar and S. Popescu, Phys. Rev. Lett. **74**, 1259 (1995).

[3] C. H. Bennett, D. P. DiVincenzo, C. A. Fuchs, T. Mor, E. Rains, P. W. Shor, J. A. Smolin, and W. K. Wootters, Phys. Rev. A **59**, 1070 (1999).

[4] G. J. Pryde, J. L. O'Brien, A. G. White, and S. D. Bartlett, Phys. Rev. Lett. **94**, 220406 (2005).

[5] J. Niset, A. Acin, U. L. Andersen, N. J. Cerf, R. García-Patrón, M. Navascues, and M. Sabuncu, quant-ph/0608215.

[6] B. M. Terhal, D. P. DiVincenzo, and D. W. Leung, Phys. Rev. Lett. **86**, 5807 (2001).

[7] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. **80**, 5239 (1998).

[8] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. **78**, 574 (1997).

[9] I. D. Ivanovic, Phys. Lett. A **123**, 257 (1987).

[10] J. A. Bergou, U. Herzog, and M. Hillery, Phys. Rev. Lett. **90**, 257901 (2003).

[11] G. Jaeger and A. Shimony, Phys. Lett. A **197**, 83 (1995).

[12] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).

[13] Y.-. X. Chen and D. Yang, Phys. Rev. A **64**, 064303 (2001).